

Comments on the Matrix Representations of Quaternions

Gisele Ducati

Departamento de Matemática, UFPR
 CP 19081, 81531-970 Curitiba (PR) Brasil
 ducati@mat.ufpr.br

February 1, 2008

Abstract Some comments are made on the matrices which serve as the basis of a quaternionic algebra. We show that these matrices are related with the quaternionic action of the imaginary units from the left and from the right.

MSC. 11R52 - 15A33.

We start by recalling the notation of the referred paper [1]. It's well known that a quaternion

$$q = a + bh + cj + dk, \quad a, b, c, d \in \mathbb{R},$$

where h, j and k denote the imaginary units of the algebra of quaternions

$$h^2 = j^2 = k^2 = hjk = -1,$$

can be written in matrix form as

$$Q = aI + bH + cJ + dK,$$

with

$$H^2 = J^2 = K^2 = HJK = -I.$$

Like pointed out in the paper [1] the matrices H, J and K can be chosen in different ways so adopting skew-symmetric signed permutation matrices with a plus one in its first row we find six signed permutation matrices which arrange themselves to form two Hamiltonian systems, given by H_1, J_1, K_1 and H_2, J_2, K_2 .

We emphasize that the first system represents the left action of the imaginary units j, h and k , respectively, and the matrices of the second system represents, respectively, the right action of the quaternionic imaginary units, h, j and k . In both cases we should multiply the units by -1 . Let $q = q_0 + hq_1 + jq_2 + kq_3$ and $p = p_0 + hp_1 + jp_2 + kp_3$ real quaternions. If we distinguish between the left and right action of the quaternionic imaginary units h, j , and k by introducing the operators L_q and R_p whose action on quaternionic functions Ψ , $\Psi : \mathbb{H} \rightarrow \mathbb{R}$, is given by

$$L_q \Psi = q \Psi \quad \text{and} \quad R_p \Psi = \Psi p, \quad (1)$$

where

$$L_h^2 = L_j^2 = L_k^2 = R_h^2 = R_j^2 = R_k^2 = L_h L_j L_k = R_k R_j R_h = -1,$$

and

$$[L_q, R_p] = L_q R_p - R_p L_q = 0, \quad (2)$$

we find the following *real* matrix representation [2, 3]

$$L_q \leftrightarrow \begin{pmatrix} q_0 & -q_1 & -q_2 & -q_3 \\ q_1 & q_0 & -q_3 & q_2 \\ q_2 & q_3 & q_0 & -q_1 \\ q_3 & -q_2 & q_1 & q_0 \end{pmatrix} \quad \text{and} \quad R_p \leftrightarrow \begin{pmatrix} p_0 & -p_1 & -p_2 & -p_3 \\ p_1 & p_0 & p_3 & -p_2 \\ p_2 & -p_3 & p_0 & p_1 \\ p_3 & p_2 & -p_1 & p_0 \end{pmatrix}. \quad (3)$$

It's interesting to note that the six matrices

$$L_h, L_j, L_k, R_h, R_j \text{ and } R_k \quad (4)$$

are the generators of the one-dimensional quaternion unitary group linear from the right. See [2, 4] for further details. In addition, it's more natural to arrange the six signed permutation matrices cited in [1] as given in (4).

Now, let us see how many triplets, obeying the quaternionic algebra, is possible to write with these elements. We should consider each element from (4) and it's conjugate (or it's negation) which give 12 elements. Obviously, fixing a imaginary unit, the second one must be chosen in a way that it does not commute with the previously fixed one, otherwise, the multiplication of the imaginary units will not square to minus one, contradicting the quaternion algebra. So, if L_h has been chosen for the first imaginary unit, the remaining possibilities are $\pm L_j$ and $\pm L_k$ and the third imaginary unit is determined by the multiplication of the first two. So we have 12 possibilities for the first imaginary unit and 4 for the second, that means, $12 \times 4 \times 1$, totalizing 48 possibilities. Of course, we can consider all automorphisms of the quaternion algebra, or alternatively, thinking in a geometric way, any quaternion can be rotated in the three dimensional space, then we should consider the transformation

$$q \rightarrow q_u = uqu^{-1}, \quad |u| = 1.$$

Note that

$$h_u^2 = j_u^2 = k_u^2 = h_u j_u k_u = -1.$$

This similarity transformation can be made with L_h, L_j, L_k, R_h, R_j and R_k . The automorphism group of the units is the $O(3)$ subgroup of $O(4)$ given by

$$\Omega' = U\Omega U^\dagger.$$

References

- [1] R. W. Farebrother, J. Groß and S. Troschke, Matrix representation of quaternions, *Lin. Alg. Appl.* **362**, 251–255 (2003).
- [2] S. De Leo and G. C. Ducati, Quaternionic groups in physics, *Int. J. Theor. Phys.* **38**, 2197–2220 (1999).
- [3] S. De Leo and G. C. Ducati, Quaternionic differential operators, *J. Math. Phys.* **42**, 2236–2265 (2001).
- [4] G. Ducati, *Operadores diferenciais quaterniônicos e aplicações em Física*, Ph.D. Thesis, IMECC-UNICAMP (2002).